



FAMILY FOOT AND ANKLE CLINIC


SPECIALISTS AND SURGEONS

At Family Foot & Ankle Clinic, we are committed to providing you with the highest quality of care and ensuring you have all the information you need to make informed decisions about your health.

If you have any questions or concerns, please don't hesitate to reach out to us. You can contact our office directly at 763-421-7300. Additionally, you can visit our website at <https://familyfootmn.com/> for more resources and information about our services.

Thank you for choosing us for your healthcare needs!

Tailor's Bunion (Bunionette)



What Is a Tailor's Bunion?

Tailor's bunion, also called a bunionette, is a prominence of the fifth metatarsal bone at the base of the little toe. The metatarsals are the five long bones of the foot. The prominence that characterizes a tailor's bunion occurs at the metatarsal head, located at the far end of the bone where it meets the toe. Tailor's bunions are not as common as bunions, which occur on the inside of the foot, but they are similar in symptoms and causes.

Why is it called a tailor's bunion? The deformity received its name centuries ago, when tailors sat cross-legged all day with the outside edge of their feet rubbing on the ground. This constant rubbing led to a painful bump at the base of the little toe.

Causes

Often a tailor's bunion is caused by an inherited faulty mechanical structure of the foot. In these cases, changes occur in the foot's bony framework, resulting in the development of an enlargement. The fifth metatarsal bone starts to protrude outward, while the little toe moves

inward. This shift creates a bump on the outside of the foot that becomes irritated whenever a shoe presses against it.

Sometimes a tailor's bunion is actually a bony spur (an outgrowth of bone) on the side of the fifth metatarsal head.

Regardless of the cause, the symptoms of a tailor's bunion are usually aggravated by wearing shoes that are too narrow in the toe, producing constant rubbing and pressure.

Symptoms

The symptoms of tailor's bunions include redness, swelling and pain at the site of the enlargement. These symptoms occur when wearing shoes that rub against the enlargement, irritating the soft tissues underneath the skin and producing inflammation.

Diagnosis

Tailor's bunion is easily diagnosed because the protrusion is visually apparent. X-rays may be ordered to help the foot and ankle surgeon determine the cause and extent of the deformity.

Nonsurgical Treatment

Treatment for tailor's bunion typically begins with nonsurgical therapies. Your foot and ankle surgeon may select one or more of the following:

- **Shoe modifications.** Choose shoes that have a wide toe box, and avoid those with pointed toes or high heels.
- **Padding.** Bunionette pads placed over the area may help reduce pain.
 - www.myfootshop.com --> Bunion Products --> Tailors Bunion Pads



- **Oral medications.** Nonsteroidal anti-inflammatory drugs (NSAIDs), such as ibuprofen, may help relieve the pain and inflammation.
- **Icing.** An ice pack may be applied to reduce pain and inflammation. Wrap the pack in a thin towel rather than placing ice directly on your skin.
- **Injection therapy.** Injections of corticosteroid may be used to treat the inflamed tissue around the joint.

- **Orthotic devices.** In some cases, custom orthotic devices may be provided by the foot and ankle surgeon.

When Is Surgery Needed?

Surgery is often considered when pain continues despite the above approaches. In selecting the procedure or combination of procedures for your case, the foot and ankle surgeon will take into consideration the extent of your deformity based on the x-ray findings, your age, your activity level and other factors. The length of the recovery period will vary, depending on the procedure or procedures performed.

[https://www.foothealthfacts.org/conditions/tailor-s-bunion-\(bunionette\)](https://www.foothealthfacts.org/conditions/tailor-s-bunion-(bunionette))