

Advanced Pediatrics of Rockland, P.C.

358 Route 202, Suite 2, Pomona, NY 10970

P: (845) 364-9800

F: (845) 364-9828

Anticipatory Guidance

2 year old - Female

You can start to transition your child to a toddler bed at this age.

You should be giving her prescription multivitamins with fluoride.

Brush her teeth twice a day with fluoride-free toothpaste since ingesting too much fluoride can cause tooth staining.

Use fragrance free, mineral sunblock for sun exposure.

Continue childproofing the house: Ensure pills and poisons are not accessible. This includes all household cleaners which should be stored in a cabinet with childproof locks on it. Clean up small pieces and coins that she can choke on or ingest. Get rid of (or pad) coffee tables or other hard/pointy objects. Pad all corners and edges of furniture to help prevent head injuries and lacerations. Secure all TV's and dressers so that they cannot fall on your child.

Do not give choking hazards such as nuts, raisins and grapes. She should continue to drink mostly water. Try not to give juice or milk since they contain empty sugar and calories without adding much nutrition. If your child is drinking milk, she should be drinking fat free (skim) milk and limit it to one cup per day. Good Karma unsweetened flax milk is a very low-calorie alternative and contains vitamin D, calcium and omega 3 fatty acids for brain development. Adding flax seed oil to food will also add omega 3 free fatty acids. Food choices should include vegetables for vitamins and minerals. It should also include protein containing foods such as meat, fish, chicken, egg whites and beans.

Continue to use "time out" and ignore temper tantrums. Remember that "time out" should be used right after your child demonstrates behavior that needs to be corrected. You should also limit its use which will increase its effectiveness. Ignoring bad behavior is still the most important part of discipline. Most kids will not sit in a "time out" chair. Simply sit her on your lap, cross your arms and hold her wrists. Set the timer on your phone for two minutes. Tell her that she is getting a "time out" and when the timer goes off, let her up. Start the "time out" as soon as she is on your lap and don't wait for her to calm down. After the "time out" is over, try to engage her when she is playing nicely in order to increase positive reinforcement.

Advanced Pediatrics of Rockland, P.C.

358 Route 202, Suite 2, Pomona, NY 10970

P: (845) 364-9800

F: (845) 364-9828

Your child should be going to bed tired but awake - This allows her to learn to self-soothe which is a very important tool for social-emotional development. She should be using a straw cup or open cup now. We strongly recommend discontinuing all bottles and sippy cups at this age if she is still using them.

Most girls will start to potty-train at this age. You should use positive reinforcement only and try not to force her to use the potty. Putting too much pressure on potty-training can lead to functional withholding and significant constipation.